

# Vis/NIR Data Quality, Cloud Detection and IR co-registration

---

Catherine Gautier and Yang Shiren  
Institute of Computational Earth System Science  
UCSB

# Overview

---

- Data Quality
  - Stripes
  - Co-alignment between Vis/NIR and AIRS
- Effect Registration between Vis/NIR and 1993 NDVI on cloud detection
- Cloud Detection
- Future Work


# Stripes

---

- Stripes caused by difference of gain and offset between 9 Vis/NIR elements
- In-between element adjustment + vicarious calibration used to remove stripes
- Remaining small stripes in different conditions
- Detailed investigation of stripes characteristics


# Gain computed over bright surface



---




# Stripes in Sunglint area

---






Ch1 max radiance=594.915





Ch2 max radiance=526.404




Ch3 max radiance=348.745


Granule 150 Box A



Ch1 max radiance=594.915






Ch2 max radiance=526.404




Ch3 max radiance=348.745

Granule 150 Box B






Ch1 max radiance=833.660





Ch2 max radiance=712.472




Ch3 max radiance=534.189


Granule 028 Box B



Ch1 max radiance=833.660



Ch2 max radiance=712.472



Ch3 max radiance=534.189

Granule 028 Box D

# Stripes Analysis: Preliminary Results


---

- Suggestion of a non-linear gain and/or BRDF effects due to scanning geometry

# Registration between Vis/NIR and IR



---

- Use transition targets to evaluate quality of Vis/IR alignment/co-registration
- Investigated E-W and N-S alignment




Gautier - Yang



13




(a) Horizontal cross section




(b) Vertical Profile



(a) Horizontal cross section






(b) Vertical Profile




Gautier - Yang

16








(a) Horizontal cross section





(b) Vertical Profile




(a) Horizontal cross section



(b) Vertical Profile



(a) Horizontal cross section



(b) Vertical Profile

# Preliminary Results

---

- Very good within accuracy of approach
- Need longer data set over same transition targets to quantitatively evaluate alignment


# Registration between Vis/NIR and 1993 NDVI



---

- 1993 NDVI used in computation of cloud threshold as a surface type (reflectance) indicator
- Preliminary analysis=>mis-alignment between NDVI 1993 and AIRS Vis/NIR NDVI
- Use transition targets to evaluate Vis/IR alignment/co-registration
- Investigate E-W and N-S alignment


# Selection of Transition Targets

---






(a) Horizontal cross section





(b) Vertical Profile  
NDV normalized for display



# Selection of Transition Targets



---





# Impact on Cloud Detection

---



# Registration between Vis/NIR and 1993 NDVI : Preliminary Results

---

- Very good N-S alignment
- 1-2 Vis/NIR pixel E-W misalignment
- Found some changes related to time difference between 2002 and 1993
- Will be developing AIRS NDVI soon


# Vis/NIR Cloud Detection

---

- Based on cloud threshold
- Cloud threshold: parameters adjusted to preliminary calibration
  - Visual assessment of cloud detection very good
- Some issues due to 1993 NDVI
- Comparison with spatial inhomogeneity approach results

# Ocean Conditions: Sun glint



---



Granule 223, July 20, 2002


# Ocean Conditions: scene selection

---




# Vis/IR cloud mask vs. spatial homo

---




# Vis/IR cloud mask vs. spatial homo

---




# Vis/IR cloud mask vs. spatial homo





# Ocean conditions: hurricane

---




# Ocean Conditions: scene selection

---




# Vis/IR cloud mask vs. spatial homo

---



T(mean) degree K	294.0 295.7 296.0	293.2 295.5 296.0	293.2 294.6 295.5
------------------	-------------------	-------------------	-------------------

T(stddev) degree K	2.66 0.40 0.27	1.24 0.51 0.36	0.79 0.66 0.40
--------------------	----------------	----------------	----------------




T(mean) degree K	293.5 294.0 294.4	292.8 291.9 294.1	287.2 286.7 291.8
------------------	-------------------	-------------------	-------------------

T(stddev) degree K	0.73 0.23 0.37	2.71 3.77 0.51	5.94 7.01 1.97
--------------------	----------------	----------------	----------------


# Vis/IR cloud mask vs. spatial homo

---




# Vis/IR cloud mask vs. spatial homo

---




# Land Conditions

---





Gau



Grav e 177, July 20, 2002

# Land Conditions: scene selection


---




Gautie

# Vis/IR cloud mask vs. spatial homo

---



# Vis/IR cloud mask vs. spatial homo




# Cloud Detection : Preliminary Results

---

- Very good visual agreement
- Impact of 1-2 Vis/NIR pixel E-W misalignment and changes related to time difference between 2002 and 1993 will be addressed by AIRS NDVI soon
- Improvements based on spatial homogeneity approach

Granule 110



Left – standard deviation of brtemp in an AMSU footprint (inhomogeneity)  
Right – inhomogeneity corrected (threshold reduction) area (black)

# Near-term Activities

---

- Cloud flag over land evaluation
- Validation of cloud detection against local meteorological data and ARM/CART data
- Improved validation of geo-location and co-registration
- Vicarious calibration

# Vicarious Calibration

---

- Surface observation-based calibration suggests some surface BRDF effects that need to be taken into account
- Development of BRDF model based on satellite observations
- Adaptation of UCSB RTM BRDF model to Railroad Valley surface conditions

# Longer-term Activities

---

- Use combined Vis/NIR and IR channels for low cloud detection
- IR Spatial Inhomogeneity vs. Vis/NIR cloud threshold
- Cloud fraction comparisons
- Vis/NIR spectral response function
- Surface radiation budget modeling
- Land surface spectral emissivity